

THE
Journal
OF
Nervous and Mental Disease

Original Articles.

ON AMAUROTIC FAMILY IDIOCY.
A DISEASE CHIEFLY OF THE GRAY MATTER OF THE CENTRAL NERVOUS SYSTEM.¹

By B. SACHS, M.D.,

PROFESSOR OF MENTAL AND NERVOUS DISEASES IN THE NEW YORK POLYCLINIC,
etc.

The clinical symptoms of amaurotic family idiocy have been described frequently and thoroughly enough. It is gratifying to note that the clinical picture as developed by the present writer a number of years ago has been generally accepted, and has been fully endorsed in the excellent articles by Hirsch, Peterson, Mohr, Patrick, Falkenheim, and a number of others. Two symptoms, in addition to those originally given, have been mentioned by Falkenheim:—"Explosive laughter" and "disturbance of deglutition." The former I have not observed, though I do not doubt its occurrence in some instances, and the latter may be taken to be either an accompaniment of the great mental and physical deterioration, or it may, at times, have the value of a "bulbar" symptom; and "bulbar" symptoms are explained easily enough, in view of our present knowledge of the morbid anatomy of the disease.

While the disease is relatively rare, the records have become so numerous that Falkenheim has been able to analyze 64 cases.* The number is increasing rapidly and mere statistical studies will

¹Read at the annual meeting of the American Neurological Association, June 5, 6, and 7, 1902.

*I have another case under observation at the present time.

soon not be needed. I would urge, however, that the clinical type as now accepted, be not disturbed because of any unusual symptoms, such as were recorded, for instance, in the cases of Claiborne, Kuh and De Bruyn. The last named author has, of his own accord, and justly so, withdrawn his case from this especial type. In the case of Claiborne the condition was complicated by a tubercle of the corpora quadrigemina, and although the characteristic change in the fundus was made out, the patient presented so many unusual features that this record should be utilized, if at all, with great caution. In Kuh's case the clinical picture was obscured and altered by the presence of an hydrocephalus.

The chief interest in the study of amaurotic family idiocy has been shifted from the clinical to the anatomical side. There is, in the main, a fortunate agreement as to the findings. There can be a difference of opinion only with respect to the interpretation to be put upon these findings. Nine years have elapsed since I have had the opportunity of studying carefully the morbid changes in a case that had been under my own observation. Since that time so many new staining methods have been introduced that I was eager to undertake this special study in pathology again, and to determine to what extent conclusions of former years would have to be changed or reinforced. For the opportunity of studying the present case I am greatly indebted to Dr. Sarah Welt, who devoted an unusual amount of time and patience to the care of this child before finally referring it to Mt. Sinai Hospital; and I am also indebted to Dr. Koplik, who, knowing my interest in the subject, was good enough to put this child entirely under my charge. As in all the other cases, the history of this one may be stated briefly:

This child, C. S., born of Hebrew parents, was two years and seven months of age at the time I first saw it at the hospital, in September, 1901. The patient's mother had had three children. The first-born is now six years of age, a fine, healthy-looking child. The second child died at the age of sixteen months. Two months before its death the mother knew that it was blind. It could never sit up straight; never uttered a sound; had no spasms, but was feverish off and on. It was thought to have tubercular meningitis, but in all probability was afflicted with this family disease. Our own patient, the third child, was born about two and one-half years after the second. During pregnancy the mother was nervous, but otherwise well; labor was normal and short. At birth and for

some time afterward the child appeared to be normal; it vomited a great deal, however, and as time passed on it was noticed that it could not sit up straight, nor did it seem to be as bright as other children. The mother thought that the child recognized her when it was somewhere between six and ten months of age, but after this period the vision gradually grew fainter and fainter. At the age of one year the child was practically blind, and since then the general weakness increased. At about this time the typical changes in the region of the macula lutea were made out. At the age of nineteen months it had a vomiting spell with convulsions. In November, 1900, it was afflicted with pneumonia, from which it recovered, but the mother states that thereafter it never moved its extremities except in convulsions. The extreme contractures in the upper and lower extremities which the child presented were developed about February, 1901. The statement was made that these came on quite suddenly after a convulsion, but of this fact we cannot be entirely certain. The child had difficulty in swallowing and was startled by the slightest noise. With the exception of increasing emaciation, the condition remained practically unchanged up to the time of death, on October 16, 1901.

The history as given above is largely based upon the careful observation of Dr. Sarah Welt, and the condition was also carefully studied by me during the four weeks preceding death. In the last two weeks of life the temperatures were often subnormal. To complete the history of the case, it may be added that the mother nursed this child to the age of eleven months. She had nursed the second child to the age of eight months, while the first healthy child was raised on sterilized milk.² While carrying the amaurotic child, the mother felt very little life, and in passing it may be noted that she was distinctly of cachectic build. From the above account of the child's condition it will readily be seen that the diagnosis of amaurotic family idiocy could be made with absolute certainty.

The autopsy was performed ten hours after death, on October 16, 1901, by Dr. Libman, Assistant Pathologist of Mt. Sinai Hospital. To him and to his chief, Dr. Mandelbaum, I am indebted for a full report. We could not get permission to remove the eyes, but with this one exception the study is complete.

Upon removal from the skull the brain weighed 784 grammes; the dura mater was rather firmly adherent to the bones.

²I make mention of these facts in order to do justice to Hirsch's theory. The facts would seem to support his theory, but for reasons to be given later on I cannot accept the same.

The cortex was so firm that the knife grated as it passed through it—a sound which I have not heard under similar conditions since the autopsy on the first subject of this disease. It was noted that the convolutions were small and that there were fissural peculiar-




Fig. 1

ities,—indications of a low type of brain, such as I recorded in the very first report made upon this disease, but such changes have not been referred to in the reports of others. In view of the possible developmental character of the disease these structural peculiarities are of some importance. The fluid in the pia was markedly increased in amount. The pituitary body was not enlarged.

Of other changes noted at the time I will only refer to an old pleurisy, and to a very advanced parenchymatous myocarditis.

The brain and spinal cord were at once put in formalin and prepared for staining by the various methods now in use. Serial sections have been laid through almost every part of the entire central nervous system. A careful study of these sections shows some deficiency in the development of the cerebral white fibers and also




Fig. 2—Section through cortex, showing changed pyramidal cells. Hematoxylin-eosin stain. Oc. 4; obj. 1-12 (Leitz).

a degeneration of the pyramidal tracts in the lateral as well as in the anterior columns of the cord. This same degeneration can be traced through the course of the pyramidal tracts in the internal capsule, crusta, pons and medulla, but much more striking than this change in the white fibers is the change in the gray matter of the central nervous system. The changes are found to be the same in the cortex of the brain, in the cranial nerve nuclei, in the anterior and posterior gray matter from the cervical to the lowest lumbar, and sacral segments of the cord. Even the sacral

ganglia show similar changes. The changes are practically those to which attention was explicitly directed by Dr. Hirsch, and which were found to be similar to those reported by me in the cortex in 1887, and were also found on a reëxamination to be contained in some of the spinal sections of the specimens examined in 1892 from my second case.

Of course these changes are brought out much more beautiful-




Fig. 3—Section through XII nucleus in medulla oblongata showing advanced changes in ganglion cells. Hematoxylin-eosin stain.

ly by the Nissl and other methods now in use. There is unquestionably some increase of the neuroglia cells throughout the central nervous system, but the chief and most remarkable changes are found in the larger ganglion cells throughout the entire extent of the central nervous axis. In the cortex as well as in the anterior gray matter of the cord there is not a normal ganglion cell to be seen. As will be noted in the specimens and illustrations here-

with presented the cell body is completely altered, the entire cell protoplasm has become disintegrated, leaving a more or less homogeneous mass, the nucleus has been shifted commonly to some part of the periphery of the cell, and instead of being distinctly differentiated from the rest of the cell-body it generally shades off by degrees into this aforesaid homogeneous mass. The nucleus is often wanting entirely, and, in some instances, the entire nucleus appears to have dropped out. The disintegration of the cell-body is so complete that in many specimens the contour alone enables one to infer that these round bodies are the remnants of ganglion




Fig. 4—Section through cervical enlargement showing cell changes in anterior gray matter. (Van Gieson stain.)

cells. These metamorphosed ganglion cells are frequently surrounded by distinct pericellular spaces and are relatively and absolutely so much enlarged that they give a very striking appearance to the cross section.

There is no doubt an entire agreement between the findings in this last case of mine and those so well reported by Dr. Hirsch before this Association a number of years ago. Curiously enough, other recent writers have not been struck by these changes in the gray matter of the brain and spinal cord. In a recent and preliminary report by Prof. Schaffer of Buda-Pest there is the account of a typical case and in the report of the post-mortem findings special stress is laid upon changes of the white fibers and nothing is said

about the gray matter, except that it is entirely normal, and yet, from the cut (Fig. 2) which accompanies his article, although it is reproduced at a low power, I suspect that these same large cell-bodies are present, such as Hirsch and myself have found and described.

The chief inference to be drawn from a study of sections laid through every part of the central nervous system in this and in other cases of amaurotic family idiocy is that the morbid process




Fig. 5—Section through a lumbar ganglion. Nissl stain showing marked cell changes.

in this disease affects primarily, or at least to a great extent, the entire gray matter of the brain and of the spinal cord. The degeneration of the white fibers of the anterior and lateral pyramidal tracts is in all probability secondary, but not nearly so great or so marked as one would expect with such advanced disease of the ganglionic tissue. It is already probable that the relative involvement of the gray and white matter may vary in different subjects of this disease.

The neurone theory has received so many severe blows of late that the somewhat anomalous condition of affairs as observed in

this case need not be especially dilated upon. The theory may have to go to the wall in part or as a whole, but the facts will stand and will call for further elucidation.

In 1887 I spoke of these changes as representing an agenetic condition pure and simple affecting the highest cortical nerve elements. At that time I was not aware of the widespread character of the morbid process. Later researches, and particularly those of Kingdon and Russell and of Hirsch, have laid stress upon the degenerative character of the disease. Much has been made of the fact that the present author has recorded the process as essentially an arrest of development, while the other writers just mentioned have insisted upon the purely degenerative and acquired character of the disease. Some writers have accepted these expressions of opinion as though they indicated a widely differing conception of amaurotic family idiocy, but the difference between the views of the authors just mentioned and of myself is not so great as it would appear at first sight.

A degenerative process does not necessarily represent an acquired or an acute affection. It is my contention now, as in former years, that degeneration will set in whenever normal growth is arrested, but life continues. A child to be afflicted with amaurotic family idiocy is born with a limited and restricted capacity for normal development. Its gray cells may do as well as those of any other child up to the age of two or three or six months, but beyond that its powers for further development will not go. To this extent, in this respect, the disease is after all a congenital affair; when normal development ceases, degeneration sets in.

Gowers has recently written a very helpful article in which he applies a similar course of reasoning to the various scleroses, and has suggested the term "abiotrophy," which I gladly accept, to indicate this arrest of normal development followed by degeneration. The various parts of the nervous system, according to Gowers, have their own vitality. "Some of them may slowly die, while the life of all the rest goes on without impairment. * * * When the failure is early it is due only to a defect which seems to be inherent, the tendency thereto inborn. We do not indeed apply the word 'death' to this slow decay of the elements; we speak of it as 'degeneration'; but the process is in many cases—perhaps in most—an essential failure of vitality." The term "abiotrophy" design-

nates a lack of vital nutrition, while the term "abiotic" would designate that which depends on defective vitality. Were we to adopt Gowers' phraseology, we might with good reason claim that amaurotic family idiocy is due chiefly to a defective vitality of the gray matter of the central nervous system.

To my mind there is an almost insuperable objection to the doctrine that so distinct a family affection is an acquired disease; yet granting the possibility of this it could only be conceivable upon the further theory of toxic or other influences acting upon the tissues of deficient embryonal potentiality. Hirsch's contention that the toxic influence may have been exerted through the mother's milk cannot be maintained for the simple reason that several of my patients have not been nursed by the mother. Assuming that amaurotic family idiocy may be an acquired disease, the toxic nature of the degenerative process has been argued from the resemblance of the cell pictures to those found in cases of poisonings of various kinds. This point deserves careful consideration, but cannot be decided unless we are certain that the cell changes as we see them represent definite changes due to definite causes.

With the refined staining methods of the day, we have learned much regarding the protoplasmic changes of the cell-body. But I ask, does acute poisoning produce cell changes distinctly different from those of chronic poisoning? Again, can we make out from inspection of the cells whether death was due to metallic poisoning, to acute infectious disease, to a condition of marasmus or what not? For the present it would be hazardous to present such a claim.

It would seem more profitable to attempt to discover other conditions in which cell changes occurred somewhat similar to those made out by Hirsch and myself.

Dr. Spiller reported the case of a boy about nine years of age, the inmate of the Pennsylvania Training School for Feeble-minded Children, who was in all probability afflicted with congenital spastic rigidity and imbecility. This child was seized with symptoms resembling those of cerebro-spinal meningitis from which he died five days after the onset. The microscopical examination of the nervous tissue, Spiller writes, showed little evidence of inflammation.

"I found cell changes in all the cell-bodies of the posterior and anterior horns of the spinal cord; in the nuclei of the cranial nerves; in the sensory as well as the motor nucleus of the fifth nerve; in the cells of Purkinje and in the parietal lobule where especially the cells of Betz were altered. These cell changes were very like those found in amaurotic family idiocy. The gross lesions of meningitis were not found." Spiller seems to interpret these cell changes to be the result of the acute process. I am tempted to question him whether they might not more properly be related to the "chronic spastic rigidity and imbecility." Are we certain enough of these cell changes to claim that they must necessarily be the product of some acute intoxication?

The findings of Rolly in a case of Little's disease (congenital spastic diplegia) are of special interest in this connection. The child was the second of twins, born in the mother's fourteenth confinement. The child was born asphyxiated, and presented all the symptoms of marked spastic rigidity (diplegia), with frequent convulsions. These symptoms continued without much change until death, at the age of about two and one-half months. Rolly lays stress on the specific history of the father, and wishes chiefly to prove the bearing of syphilis upon the etiology of Little's disease. For my present purposes I am interested in his cell findings, Nissl himself having examined Rolly's specimens taken from the spinal cord and from the brain. In the lumbar portion of the cord the author states: "Side by side with cells which present a tolerably normal appearance are others which exhibit a granular disintegration of the Nissl bodies surrounding the nucleus, while the peripheral portions of the cell body remain practically normal." Similar changes were found in the dorsal portion of the cord. In the cervical portion "there are fewer normal cells than in the other divisions of the cord. As a rule, the nucleus cannot be distinctly differentiated from the cell protoplasm. More marked still are the degenerative changes in the brain. There is an enormous and diffuse proliferation of the neuroglia. Some ganglion cells have lost their contours and are more or less shapeless masses, which shade off into the surrounding tissue. Some ganglion cells are so thoroughly disintegrated that one sees practically only lacunæ, which, by their shape, remind one of the ganglion cell form. Small vacuoles were also seen to be present in some of the ganglion cells."

All of these changes bear the closest resemblance to the findings in amaurotic family idiocy. Had this child lived to the age of the amaurotic children previously described, the change would no doubt have been still more striking.

I would also call attention to similar findings in the cases of "generalized rigidity" as described by Collier,³ and in a somewhat similar case reported by Mya and Levi. Whether these latter cases are described as spastic diplegias or as cases of "spastic pseudo-bulbar paralysis," according to Peritz, the point that is of special interest to us is that very similar anatomical changes have been found in the brains of children who had been afflicted with varying forms of cerebral disease.

Peritz is inclined to the opinion that I have made a special effort to establish amaurotic family idiocy as an entirely independent disease. In this he is mistaken, for I have contended for nothing more than that this family disease represents a distinct and easily recognizable clinical type. This is evident from the fact that in this city not only neurologists, but pediatricists and general practitioners have learned to diagnosticate amaurotic family idiocy with surprising ease on clinical grounds. The disease can be differentiated easily from the other forms of idiocy and from the various types of infantile cerebral palsies. In his earliest publications on this subject the present writer endeavored to show the relationship between amaurotic family idiocy and other congenital cerebral conditions, but he considered it necessary to proceed cautiously before arriving at any definite conclusion. At the present time, however, in view of the researches of Hirsch, Rolly, Spiller, Collier, Peritz, Sachs and others, it may definitely be stated that there is a close anatomical relationship between amaurotic family idiocy and other cerebral diseases of childhood which are dependent on an arrest of, or at least a disturbance in, the normal development of the central nervous system.

I will not as yet claim that the morbid process in all of these children is identical, but it seems to me that the only just inference to be drawn is this, that in amaurotic family idiocy the entire central gray matter is the seat of an intense degeneration, and that somewhat similar changes are found in conditions more or less

³Collier calls attention to the resemblance between his anatomical findings and those of amaurotic family idiocy.

closely allied to this special type of family disease. But even if closely allied to other affections, the clinical picture of the type is so distinct that we can well afford to consider it a special clinical form of family disease. Why children of one race should be afflicted so much more often than those of others, when the allied conditions show no such preference, remains as great a puzzle as ever.

We have not yet reached the end of this discussion, and I trust that with every new case carefully reported and carefully studied further light will be thrown upon a disease that appears to me to be of increasingly greater interest. Above all things, I would have future writers on this subject pay the closest attention to the changes in the gray matter of the brain and of the spinal cord.

REFERENCES.

- Claiborne: Archives of Pediatrics, 1900, Vol. X, p. 3.
 Collier: Brain, 1899, Vol. XXII, p. 373.
 De Bruyn: Weekblad van het Med. Tidischr. v. Geneesk, 1900, Bd. II, p. 499 (quoted by Falkenheim).
 Falkenheim: Jahrbuch f. Kinderheilkunde N. F. LIV, H. 2, 1901 (contains very complete literature).
 Gowers: Lancet, April 12, 1902.
 Hirsch, William: JOURNAL OF NERVOUS AND MENTAL DISEASE, 1898, Vol. XXV, p. 538.
 Kingdon and Russell: Medico-Chirurgical Transactions, 1897, Vol. LXXX.
 Kuh: JOURNAL OF NERVOUS AND MENTAL DISEASE, Vol. XXVII, 1900, p. 268.
 Mohr: Arch. f. Augenheilkunde, Bd. 43, Heft 2, p. 285.
 Mya und Levi: Riv. de Path. nervosa e. mentale, 1896 (quoted by Peritz).
 Patrick: JOURNAL OF NERVOUS AND MENTAL DISEASE, Vol. XXVII, 1900, p. 265.
 Peritz: Die Pseudo-bulbaer und Bulbaerparalysen des Kindesalters; S. Karger, Berlin, 1902.
 Peterson: JOURNAL OF NERVOUS AND MENTAL DISEASE, 1898, Vol. XXV, p. 529.
 Rolly: Zeitschr. f. Nervenheilkunde, Bd. XX, p. 152.
 Sachs, B.: "On Arrested Cerebral Development," etc. JOURNAL OF NERVOUS AND MENTAL DISEASE, New York, 1887, p. 541; 1892, p. 603. New York Medical Journal, 1896, May 30. Volkmann'scher Vortrag, N. F. No. 46, 47 (Die Hirnlaehmungen der Kinder), p. 34.
 Shaffer: Wiener klinische Rundschau, No. 16, 1902.
 Spiller: "A Case with the Symptoms of Cerebro-Spinal Meningitis," etc. JOURNAL OF NERVOUS AND MENTAL DISEASE, 1901, p. 140.